

Trigonometric Degeneration and Orbifold Wess-Zumino-Witten Model. I

Takashi Takebe
Department of Mathematics
Ochanomizu University
Otsuka 2-1-1, Bunkyo-ku
Tokyo, 112-8610, Japan

February 2003; Revised: March 2003

Abstract

Trigonometric degeneration of the Baxter-Belavin elliptic r matrix is described by the degeneration of the twisted WZW model on elliptic curves. The spaces of conformal blocks and conformal coinvariants of the degenerate model are factorised into those of the orbifold WZW model.

1 Introduction

The Baxter-Belavin elliptic classical r matrix is a quasi-classical limit of the famous Baxter's elliptic R matrix.[1] It has a unique position in the classification of the classical r matrices by Belavin and Drinfeld.[2]

Etingof introduced the elliptic Knizhnik-Zamolodchikov (KZ) equation as a system of differential equations satisfied by the twisted trace of product of vertex operators (intertwining operators of representations) of affine Lie algebras $\widehat{sl}_N$. [3] This system is formally obtained by replacing the rational r matrix in the ordinary KZ equation by the elliptic one. Kuroki and the

author constructed the *twisted Wess-Zumino-Witten (WZW) model* on elliptic curves and gave to Etingof's equations a geometric interpretation as flat connections.[4]

As is well-known, the elliptic r matrix degenerates to the trigonometric r matrix when the elliptic modulus τ tends to $i\infty$. Hence it is natural to expect that the twisted WZW model becomes a degenerate model which is related to the trigonometric r matrix when the elliptic curve degenerates to a singular curve.

The purpose of this and the subsequent papers is to show that this is indeed the case and to analyse the degenerate WZW model. In the present paper we examine the degenerate WZW model and prove the factorisation theorem of Tsuchiya-Ueno-Yamada type.[5] The spaces of conformal blocks and conformal coinvariants of the twisted WZW model over the singular curve factorise into those of the orbifold WZW model, which has been intensively studied in other context. See, e.g., Ref. [6]. Actually, though based on a different standpoint, Etingof's work[3] could be considered as a precursor to them.

In the forthcoming papers we study the behaviour of the sheaves of conformal blocks and conformal coinvariants of the twisted WZW model around the boundary of the moduli of elliptic curves, especially the factorisation property of these sheaves. The integrable systems arising from this WZW model, i.e., the trigonometric Gaudin model and the trigonometric KZ equations, shall be also analysed.

In this work, we use the formulation of WZW models in terms of the affine Lie algebras. The reader can translate it into the formulation by the vertex operator algebras[7, 8, 9] without difficulties.

This paper is organised as follows: After reviewing the twisted WZW model on elliptic curves in §2, we define the trigonometric twisted WZW model and corresponding WZW model on the orbifold in §3. The space of conformal coinvariants of the trigonometric twisted WZW model is decomposed into the “direct integral” of the spaces of conformal coinvariants of the orbifold WZW model. This general factorisation theorem is proved in §4. If certain conditions are imposed on the modules inserted to the degenerate curve, a refined factorisation theorem can be proved. This is done in §5. The simplest non-trivial example of the factorisation is given in §6. The last section §7 summarises the results and lists several problems.

Notations

We use the following notations besides other ordinary conventions in mathematics.

- N, L : fixed integers. $N \geq 2$ will be the rank of the Lie algebra and $L \geq 1$ will be the number of the marked points on a curve.

- Let X be an algebraic variety, $\mathcal{F}$ be a sheaf on X and P be a point on X .

$\mathcal{O}_X, \Omega_X^1, \mathcal{F}_P, \mathfrak{m}_P$: the structure sheaf of X , the sheaf of holomorphic 1-forms on X , the stalk of $\mathcal{F}$ at P , the maximal ideal of the local ring $\mathcal{O}_{X,P}$, respectively.

When $\mathcal{F}$ is an $\mathcal{O}_X$ -module;

$\mathcal{F}|_P := \mathcal{F}_P / \mathfrak{m}_P \mathcal{F}_P, \mathcal{F}_P^\wedge$: fibre of $\mathcal{F}$ at P , $\mathfrak{m}_P$ -adic completion of $\mathcal{F}_P$, respectively.

- We shall use the same symbol for a vector bundle and for a locally free coherent $\mathcal{O}_X$ -module consisting of its local holomorphic sections.

2 Review of twisted WZW model on elliptic curves

In this section we briefly review the twisted WZW model on elliptic curves. In the next section the degeneration of this model, the trigonometric twisted WZW model, shall be given. We follow Ref. [4] but make slight changes of normalisation and notations because we need to use the results of Etingof[3] later.

We fix an invariant inner product of $\mathfrak{g} = \mathfrak{sl}_N(\mathbb{C})$ by

$$(A|B) := \text{tr}(AB) \quad \text{for } A, B \in \mathfrak{g}. \quad (1)$$

Define matrices β and γ by

$$\beta := \begin{pmatrix} 0 & 1 & & 0 \\ & 0 & \ddots & \\ & & \ddots & 1 \\ 1 & & & 0 \end{pmatrix}, \quad \gamma := \begin{pmatrix} 1 & & & 0 \\ & \varepsilon^{-1} & & \\ & & \ddots & \\ 0 & & & \varepsilon^{1-N} \end{pmatrix}, \quad (2)$$

where $\varepsilon = \exp(2\pi i/N)$. Then we have $\beta^N = \gamma^N = 1$ and $\gamma\beta = \varepsilon\beta\gamma$.

Let $E = E_\tau$ be the elliptic curve with modulus τ : $E_\tau := \mathbb{C}/\mathbb{Z} + \tau\mathbb{Z}$. We define the group bundle G^{tw} with fibre $G = SL_N(\mathbb{C})$ and its associated Lie algebra bundle $\mathfrak{g}^{\text{tw}}$ with fibre $\mathfrak{g} = sl_N(\mathbb{C})$ over E by

$$G^{\text{tw}} := (\mathbb{C} \times G)/\sim, \quad (3)$$

$$\mathfrak{g}^{\text{tw}} := (\mathbb{C} \times \mathfrak{g})/\approx, \quad (4)$$

where the equivalence relations $\sim$ and $\approx$ are defined by

$$(z, g) \sim (z + 1, \gamma g \gamma^{-1}) \sim (z + \tau, \beta g \beta^{-1}), \quad (5)$$

$$(z, A) \approx (z + 1, \text{Ad } \gamma(A)) \approx (z + \tau, \text{Ad } \beta(A)). \quad (6)$$

(Because $\gamma\beta = \varepsilon\beta\gamma$, the group bundle G^{tw} is *not* a principal G -bundle.¹) Let $J_{ab} = \beta^a \gamma^{-b}$, which satisfies

$$\text{Ad } \gamma(J_{ab}) = \varepsilon^a J_{ab}, \quad \text{Ad } \beta(J_{ab}) = \varepsilon^b J_{ab}. \quad (7)$$

Global meromorphic sections of $\mathfrak{g}^{\text{tw}}$ are linear combinations of $J_{ab}f(z)$ ($a, b = 0, \dots, N-1$, $(a, b) \neq (0, 0)$), where $f(z)$ is a meromorphic function with quasi-periodicity,

$$f(z + 1) = \varepsilon^a f(z), \quad f(z + \tau) = \varepsilon^b f(z). \quad (8)$$

The twisted Lie algebra bundle $\mathfrak{g}^{\text{tw}}$ has a natural connection, $\nabla_{d/dz} = d/dz$. We define the invariant $\mathcal{O}_E$ -inner product on $\mathfrak{g}^{\text{tw}}$ (regarded as an $\mathcal{O}_E$ -Lie algebra sheaf) by

$$(A|B) := \frac{1}{2N} \text{tr}_{\mathfrak{g}^{\text{tw}}}(\text{ad } A \text{ ad } B) \in \mathcal{O}_E \quad (9)$$

for sections A, B of $\mathfrak{g}^{\text{tw}}$, where the symbol ad denotes the adjoint representation of the $\mathcal{O}_E$ -Lie algebra $\mathfrak{g}^{\text{tw}}$. This inner product is invariant under the translations with respect to the connection $\nabla : \mathfrak{g}^{\text{tw}} \rightarrow \mathfrak{g}^{\text{tw}} \otimes_{\mathcal{O}_E} \Omega_E^1$:

$$d(A|B) = (\nabla A|B) + (A|\nabla B) \in \Omega_E^1 \quad \text{for } A, B \in \mathfrak{g}^{\text{tw}}. \quad (10)$$

¹The bundle G^{tw} is a PGL_N principal bundle. Hurtubise and Markman (for example, see Ref. [10]) take this viewpoint, but we prefer to regard it as a group scheme over the elliptic curve by the reason we stated in §0 of Ref. [4].

Under the trivialisation of $\mathfrak{g}^{\text{tw}}$ defined by the construction (4), the connection ∇ and the inner product $(\cdot|\cdot)$ on $\mathfrak{g}^{\text{tw}}$ coincide with the exterior derivative by z and the inner product defined by (1) respectively.

For each point P on E , we define Lie algebras,

$$\mathfrak{g}^P := (\mathfrak{g}^{\text{tw}} \otimes_{\mathcal{O}_E} \mathcal{K}_E)_P^\wedge, \quad (11)$$

where $\mathcal{K}_E$ is the sheaf of meromorphic functions on E and $(\cdot)_P^\wedge$ means the completion of the stalk at P with respect to the maximal ideal $\mathfrak{m}_P$ of $\mathcal{O}_P$. In other words, it is isomorphic to the loop Lie algebra $\mathfrak{g}((z - z_0))$, where z_0 is the coordinate of P , but the isomorphism depends on choices of the coordinate z and the trivialisation of $\mathfrak{g}^{\text{tw}}$. The subspace

$$\mathfrak{g}_+^P := (\mathfrak{g}^{\text{tw}})_P^\wedge \cong \mathfrak{g}[[z - z_0]] \quad (12)$$

of $\mathfrak{g}^P$ is a Lie subalgebra.

Let us fix mutually distinct points $Q_1, \dots, Q_L$ on E whose coordinates are $z = z_1, \dots, z_L$ and put $D := \{Q_1, \dots, Q_L\}$. We shall also regard D as a divisor on E (i.e., $D = Q_1 + \dots + Q_L$). The Lie algebra $\mathfrak{g}^D := \bigoplus_{i=1}^L \mathfrak{g}^{Q_i}$ has a 2-cocycle defined by

$$c_a(A, B) := \sum_{i=1}^L c_{a,i}(A_i, B_i), \quad c_{a,i}(A_i, B_i) := \text{Res}_{Q_i}(\nabla A_i | B_i), \quad (13)$$

where $A = (A_i)_{i=1}^L, B = (B_i)_{i=1}^L \in \mathfrak{g}^D$ and Res_{Q_i} is the residue at Q_i . (The symbol “ c_a ” stands for “Cocycle defining the Affine Lie algebra”.) We denote the central extension of $\mathfrak{g}^D$ with respect to this cocycle by $\hat{\mathfrak{g}}^D$:

$$\hat{\mathfrak{g}}^D := \mathfrak{g}^D \oplus \mathbb{C}\hat{k},$$

where $\hat{k}$ is a central element. Explicitly the bracket of $\hat{\mathfrak{g}}^D$ is represented as

$$[A, B] = ([A_i, B_i]^\circ)_{i=1}^L \oplus c_a(A, B)\hat{k} \quad \text{for } A, B \in \mathfrak{g}^D, \quad (14)$$

where $[A_i, B_i]^\circ$ are the natural bracket in $\mathfrak{g}^{Q_i}$. The Lie algebra $\hat{\mathfrak{g}}^P$ for a point P is nothing but the affine Lie algebra $\hat{\mathfrak{g}}$ of type $A_{N-1}^{(1)}$ (a central extension of the loop algebra $\mathfrak{g}((t - z)) = \mathfrak{sl}_N(\mathbb{C}((t - z)))$).

The affine Lie algebra $\hat{\mathfrak{g}}^{Q_i}$ can be regarded as a subalgebra of $\hat{\mathfrak{g}}^D$. The subalgebra $\mathfrak{g}_+^{Q_i}$ of $\mathfrak{g}^{Q_i}$ (cf. (12)) can be also regarded as a subalgebra of $\hat{\mathfrak{g}}^{Q_i}$ and $\hat{\mathfrak{g}}^D$.

Let $\mathfrak{g}_{\text{out}}$ be the space of global meromorphic sections of $\mathfrak{g}^{\text{tw}}$ which are holomorphic on E except at D :

$$\mathfrak{g}_{\text{out}} := \Gamma(E, \mathfrak{g}^{\text{tw}}(*D)). \quad (15)$$

There is a natural linear map from $\mathfrak{g}_{\text{out}}$ into $\mathfrak{g}^D$ which maps a meromorphic section of $\mathfrak{g}^{\text{tw}}$ to its germ at Q_i 's. The residue theorem implies that this linear map is extended to a Lie algebra injection from $\mathfrak{g}_{\text{out}}$ into $\hat{\mathfrak{g}}^D$, which allows us to regard $\mathfrak{g}_{\text{out}}$ as a subalgebra of $\hat{\mathfrak{g}}^D$.

Definition 2.1. The space of *conformal coinvariants* $\text{CC}_E(M)$ and that of *conformal blocks* $\text{CB}_E(M)$ over E associated to $\hat{\mathfrak{g}}^{Q_i}$ -modules M_i with the same level $\hat{k} = k$ are defined to be the space of coinvariants of $M := \bigotimes_{i=1}^L M_i$ with respect to $\mathfrak{g}_{\text{out}}$ and its dual:

$$\text{CC}_E(M) := M/\mathfrak{g}_{\text{out}}M, \quad \text{CB}_E(M) := \text{Hom}_{\mathbb{C}}(M/\mathfrak{g}_{\text{out}}M, \mathbb{C}). \quad (16)$$

(In Ref. [5], $\text{CC}_E(M)$ and $\text{CB}_E(M)$ are called the space of *covacua* and that of *vacua* respectively.)

3 Trigonometric twisted WZW model and orbifold WZW model

In this section we define the degeneration of the twisted WZW model in the previous section, §2, i.e., the trigonometric twisted WZW model on a singular “curve”. (Exactly speaking, the model is defined on a “stack”, whatever it is.) As is shown in the sections coming later, §4 and §5, if we desingularise the “curve”, the spaces of conformal coinvariants and conformal blocks are factorised into those of the orbifold WZW model, the definition of which is also given in this section.

When the modulus τ of the elliptic curve E_τ tends to $i\infty$, one of the cycle of the torus is pinched and we obtain a singular curve of the “croissant bread” shape. Such an object is constructed simply by identifying the north pole (∞) and the south pole (0) of the Riemann sphere $\mathbb{P}^1(\mathbb{C})$.

Unfortunately, the twisted bundle $\mathfrak{g}^{\text{tw}}$ over E_τ does not simply degenerates to a bundle over such a singular curve. We need to take several steps:

1. Lift the bundle $\mathfrak{g}^{\text{tw}}$ to a covering space of E_τ where the pull-back of $\mathfrak{g}^{\text{tw}}$ become trivial. The covering is N^2 -fold and the covering space is an elliptic curve isomorphic to E_τ .




Figure 1: Degeneration of an elliptic curve and its N^2 -covering.

2. Take such degeneration of the covering elliptic curve that it becomes the N^2 -fold covering of the singular curve mentioned above. This covering space consists of N copies of $\mathbb{P}^1$ connected to form a ring.
3. The “degenerate bundle” $\mathfrak{g}^{\text{tw}}$ over the singular curve is understood as the trivial bundle over the singular covering space constructed above with the cyclic group action together.

A “manifold with local finite group action” is called an *orbifold*, but in our case the underlying space has a singularity, hence it should be regarded as a *stack* in algebraic geometry. It is not our task (and certainly beyond the reach of the author’s ability) to construct a general theory of the WZW model over stacks. We shall define the twisted WZW model over the degeneration of the elliptic curve below by down-to-earth terminology only.

Let us denote the standard coordinate of $\mathbb{P}^1(\mathbb{C})$ by t . The cyclic group $C_N = \mathbb{Z}/N\mathbb{Z}$ acts as $t \mapsto \varepsilon^a t$ ($a \in C_N$) and the quotient $E_{\text{orb}} = \mathbb{P}^1/C_N$ is an ordinary orbifold. The degenerate elliptic curve E_{trig} is defined by identifying 0 and ∞ (i.e., the fixed points of the C_N action) of E_{orb} .

Not defining what the twisted bundle $\mathfrak{g}^{\text{trig}}$ over E_{trig} and the twisted bundle $\mathfrak{g}^{\text{orb}}$ over E_{orb} are, we directly define what their meromorphic sections

are in the following way. (In fact, specifying sections is equivalent to defining bundles in algebraic geometry.)

Definition 3.1. A $\mathfrak{g}$ -valued meromorphic function $f(t)$ on $\mathbb{P}^1$ is a meromorphic section of $\mathfrak{g}^{\text{orb}}$ if and only if it satisfies

$$f(\varepsilon t) = \text{Ad } \gamma(f(t)). \quad (17)$$

The same $f(t)$ is a section of $\mathfrak{g}^{\text{trig}}$ if and only if it satisfies

$$f(\infty) = \text{Ad } \beta(f(0)), \quad (18)$$

in addition to (17).

The relations (18) and (17) are degenerate form of (6).

Hereafter we fix mutually distinct points $Q_1, \dots, Q_L$ on the orbifold E_{orb} with coordinates $t = t_1, \dots, t_L$ (modulo C_N -action). The algebras $\mathfrak{g}^{\text{trig}, Q_i}$, $\mathfrak{g}_+^{\text{trig}, Q_i}$, $\mathfrak{g}^{\text{orb}, Q_i}$ and $\mathfrak{g}_+^{\text{orb}, Q_i}$ are defined from $\mathfrak{g}^{\text{trig}}$ and $\mathfrak{g}^{\text{orb}}$ in the same way as $\mathfrak{g}^{Q_i}$ and $\mathfrak{g}_+^{Q_i}$ for the elliptic curve, (11) and (12). When $Q_i \neq 0, \infty$ ($i = 1, \dots, L$), they are isomorphic to $\mathfrak{g}^{Q_i}$ and $\mathfrak{g}_+^{Q_i}$ since the definition is local.

For the orbifold WZW model, we need similar Lie algebras $\mathfrak{g}^{(0)}$, $\mathfrak{g}_+^{(0)}$, $\mathfrak{g}^{(\infty)}$, $\mathfrak{g}_+^{(\infty)}$ sitting at the singular points, 0 and ∞ . Formally their definition is the same as (11) and (12) with P replaced by 0 and ∞ and E by E_{orb} . The mode expansion of $\mathfrak{g}^{(0)}$ has a different form from the usual loop algebra, namely,

$$\mathfrak{g}^{(0)} = \bigoplus_{\substack{a,b=0,\dots,N-1 \\ (a,b) \neq (0,0)}} \bigoplus_{m \in \mathbb{Z}} \mathbb{C} J_{a,b} \otimes t^{a+mN}. \quad (19)$$

The algebra $\mathfrak{g}^{(\infty)}$ has the same form.

The central extension of $\mathfrak{g}^{Q_i}$, $\hat{\mathfrak{g}}^{(Q_i)}$, is defined in the same way as in §2 by using the cocycle (13). The central extension of $\mathfrak{g}^{(0)}$ and $\mathfrak{g}^{(\infty)}$ are defined essentially in the same manner. Difference is that we have to change the normalisation of the central element $\hat{k}$:

$$[A, B] = [A, B]^\circ \oplus c_{a,0}(A, B) \hat{k}, \quad c_{a,0}(A, B) := \frac{1}{N} \text{Res}_{t=0}(\nabla A | B) \quad (20)$$

for $A, B \in \mathfrak{g}^{(0)}$, where $[A, B]^\circ$ is the natural loop-algebra bracket in $\mathfrak{g}^{(0)}$. The origin of the factor $1/N$ shall be explained soon later. The definition of $\hat{\mathfrak{g}}^{(\infty)}$ is exactly the same. We often denote $\mathfrak{g}^{(0)}$ and $\mathfrak{g}^{(\infty)}$ by $\mathfrak{g}_\gamma^{(0)}$ and $\mathfrak{g}_\gamma^{(\infty)}$, following

§1 of Ref. [3], where $\hat{\mathfrak{g}}_\gamma$ is shown to be isomorphic to the ordinary affine Lie algebra $\hat{\mathfrak{g}}$. The suffix γ is put to emphasise the twisting of the algebra by γ , (17).

Let D be the divisor $Q_1 + \cdots + Q_L$. The definition of the loop algebra $\mathfrak{g}^D$ and its central extension does not change so far as D does not contain 0 and ∞ . When D contains 0 or ∞ or both, then the change of the normalisation (20) of the cocycle at 0 and ∞ should be taken into account. (When we talk about E_{trig} or $\mathfrak{g}^{\text{trig}}$, it is always assumed that D contains neither 0 nor ∞ .)

Instead of the subalgebra $\mathfrak{g}_{\text{out}}$, (15), we define the following subalgebras of $\hat{\mathfrak{g}}^D$:

$$\mathfrak{g}_{\text{out}}^{\text{trig}} := \{f(t) : \text{global meromorphic section of } \mathfrak{g}^{\text{trig}} \text{ with poles in } D\}. \quad (21)$$

$$\mathfrak{g}_{\text{out}}^{\text{orb}} := \{f(t) : \text{global meromorphic section of } \mathfrak{g}^{\text{orb}} \text{ with poles in } D\}, \quad (22)$$

Their Lie algebra structures are defined by the pointwise Lie bracket of $\mathfrak{g}$ -valued functions. It is not trivial that they are subalgebras of $\hat{\mathfrak{g}}^D$, since the residue theorem cannot be directly applied. For $f(t), g(t) \in \mathfrak{g}_{\text{out}}^{\text{trig}}$, it is easy to see that

$$c_a(f, g) = \sum_{i=1}^L \text{Res}_{Q_i}(\nabla f(t)|g(t)) = \frac{1}{N} \sum_{a=0}^{N-1} \sum_{i=1}^L \text{Res}_{\varepsilon^a Q_i}(\nabla f(t)|g(t)) = 0,$$

where $\varepsilon^a Q_i$ denotes the action of $a \in C_N$ on the point Q_i . Since $\{\varepsilon^a Q_i \mid a \in C_N, i = 1, \dots, L\}$ contains all poles of $(\nabla f(t)|g(t))$ on $\mathbb{P}^1$, we can apply the residue theorem here. This proves that $\mathfrak{g}_{\text{out}}^{\text{trig}}$ defined by (21) is a Lie subalgebra of $\hat{\mathfrak{g}}^D$. In the case of $\mathfrak{g}_{\text{out}}^{\text{orb}}$, when D does not contain 0 nor ∞ , the proof is the same. When D contains, e.g., 0, the central extension should be k times

$$\begin{aligned} c_a(f, g) &= c_{a,0}(f(t), g(t)) + \sum_{i=1}^L c_{a,i}(f(t)|g(t)) \\ &= \frac{1}{N} \left(\text{Res}_0(\nabla f(t)|g(t)) + \sum_{a=0}^{N-1} \sum_{i=1}^L \text{Res}_{\varepsilon^a Q_i}(\nabla f(t)|g(t)) \right) = 0. \end{aligned}$$

The last equality is the consequence of the residue theorem applied to $(\nabla f(t)|g(t))$ over $\mathbb{P}^1$. Hence, also in this case $\mathfrak{g}_{\text{out}}^{\text{orb}}$ is a subalgebra of $\hat{\mathfrak{g}}^D$. Other cases (when D contains ∞) are proved in the same way. This is the reason why we have to put $1/N$ in (20), which appeared first in Ref. [3], Lemma 1.1, where Etingof

wrote down explicit formulae of the isomorphism of $\hat{\mathfrak{g}}_\gamma$ and the ordinary affine Lie algebra $\hat{\mathfrak{g}}$.

We are now ready to define the spaces of conformal coinvariants and conformal blocks for E_{trig} and E_{orb} .

Definition 3.2. (i) The space of *conformal coinvariants* $\text{CC}_{\text{trig}}(M)$ and that of *conformal blocks* $\text{CB}_{\text{trig}}(M)$ over E_{trig} associated to $\hat{\mathfrak{g}}^{Q_i}$ -modules M_i with the same level $\hat{k} = k$ are defined to be the space of coinvariants of $M := \bigotimes_{i=1}^L M_i$ with respect to $\mathfrak{g}_{\text{out}}^{\text{trig}}$ and its dual:

$$\text{CC}_{\text{trig}}(M) := M/\mathfrak{g}_{\text{out}}^{\text{trig}}M, \quad \text{CB}_{\text{trig}}(M) := \text{Hom}_{\mathbb{C}}(M/\mathfrak{g}_{\text{out}}^{\text{trig}}M, \mathbb{C}). \quad (23)$$

(ii) The space of *conformal coinvariants* $\text{CC}_{\text{orb}}(M)$ and that of *conformal blocks* $\text{CB}_{\text{orb}}(M)$ over E_{orb} associated to $\hat{\mathfrak{g}}^{Q_i}$ -modules M_i with the same level $\hat{k} = k$ are defined to be the space of coinvariants of $M := \bigotimes_{i=1}^L M_i$ with respect to $\mathfrak{g}_{\text{out}}^{\text{orb}}$ and its dual:

$$\text{CC}_{\text{orb}}(M) := M/\mathfrak{g}_{\text{out}}^{\text{orb}}M, \quad \text{CB}_{\text{orb}}(M) := \text{Hom}_{\mathbb{C}}(M/\mathfrak{g}_{\text{out}}^{\text{orb}}M, \mathbb{C}). \quad (24)$$

In this paper we consider only *smooth modules* of affine Lie algebras $\hat{\mathfrak{g}}^{Q_i}$, $\hat{\mathfrak{g}}^{(0)}$, etc.: Let $\hat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[\xi, \xi^{-1}] \oplus \mathbb{C}\hat{k}$ be an affine Lie algebra. (We identify $\hat{\mathfrak{g}}^{Q_i}$, $\hat{\mathfrak{g}}^{(0)}$ and so on with this algebra by means of suitable isomorphisms.) A $\hat{\mathfrak{g}}$ -module M is called *smooth* if for any $v \in M$, there exists $m \geq 0$ such that, for $A_1, \dots, A_\nu \in \mathfrak{g}$, $m_1, \dots, m_\nu \geq 0$, and $\nu = 0, 1, 2, \dots$,

$$(A_1 \otimes \xi^{m_1}) \cdots (A_\nu \otimes \xi^{m_\nu})v_i = 0 \quad \text{if } m_1 + \cdots + m_\nu \geq m, \quad (25)$$

where ξ is the loop parameter of $\hat{\mathfrak{g}} = \mathfrak{g}((\xi)) \oplus \mathbb{C}\hat{k}$. Not only Laurent polynomials but also Laurent series in $\mathfrak{g} \otimes \mathbb{C}[\xi^{-1}][[\xi]]$ can act on smooth modules. Therefore, if M_i ($i = 1, \dots, L$) are smooth $\hat{\mathfrak{g}}^{Q_i}$ modules, the actions of $\mathfrak{g}_{\text{out}}^{\text{trig}}$ and $\mathfrak{g}_{\text{out}}^{\text{orb}}$ on $M_1 \otimes \cdots \otimes M_L$ are well-defined.

Concrete examples of spaces of conformal coinvariants shall be given in §6. The relation of CC_{trig} and CC_{orb} is our main subject in this paper and studied in the next sections, §§4–5. How CC_E for elliptic curves degenerates to these spaces at the boundary of the moduli space of the elliptic curves shall be clear in the forthcoming papers.

4 Factorisation (1)

In this section the general factorisation theorem of the space of conformal coinvariants CC_{trig} is shown. The result is not so sharp as that in the next section, but the idea of the factorisation becomes clear.

To state the theorem, we need the notion of the Verma module of $\hat{\mathfrak{g}}_\gamma$, i.e., $\hat{\mathfrak{g}}^{(0)}$ and $\hat{\mathfrak{g}}^{(\infty)}$. Recall that $\hat{\mathfrak{g}}^{(0)}$ has a natural triangular decomposition:

$$\hat{\mathfrak{g}}^{(0)} = \mathfrak{n}_+^{(0)} \oplus \mathfrak{h}^{(0)} \oplus \mathfrak{n}_-^{(0)} \oplus \mathbb{C}\hat{k}, \quad (26)$$

where $\mathfrak{n}_+^{(0)}$ consists of $\mathfrak{g}$ -valued power series in t with strictly positive powers while $\mathfrak{n}_-^{(0)}$ consists of series in t with negative powers. (See (19).) $\mathfrak{h}^{(0)}$ is the same as the Cartan subalgebra of $\mathfrak{g}$, which is spanned by $J_{0,b}$ ($b = 1, \dots, N-1$). Since $\mathfrak{h}^{(0)}$ is canonically identified with the Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$, we often drop the superscript (0). The algebra $\hat{\mathfrak{g}}^{(\infty)}$ has the same decomposition but with opposite powers of t : $\mathfrak{n}_+^{(\infty)}$ is the subspace of $\mathfrak{g}^{(\infty)}$ with negative powers of t and $\mathfrak{n}_-^{(\infty)}$ is that with negative powers of t , because $\mathfrak{n}_+^{(\infty)}$ is the space of $\mathfrak{g}$ valued functions which vanish at ∞ . To avoid repeating the same statements for $\hat{\mathfrak{g}}^{(0)}$ and $\hat{\mathfrak{g}}^{(\infty)}$, we identify them in the canonical way and denote the above decomposition as

$$\hat{\mathfrak{g}}_\gamma = \mathfrak{n}_{\gamma,+} \oplus \mathfrak{h}_\gamma \oplus \mathfrak{n}_{\gamma,-} \oplus \mathbb{C}\hat{k}, \quad (27)$$

and define $\hat{\mathfrak{b}}_\gamma := \mathfrak{n}_{\gamma,+} \oplus \mathfrak{h}_\gamma \oplus \mathbb{C}\hat{k}$.

Definition 4.1. (i) Let λ is an element of $\mathfrak{h}_\gamma^* = \text{Hom}_{\mathbb{C}}(\mathfrak{h}_\gamma, \mathbb{C})$ and k is an arbitrary complex number. The *Verma* module of $\hat{\mathfrak{g}}_\gamma$ of highest weight λ and level k is the $\hat{\mathfrak{g}}_\gamma$ -module

$$M_\lambda := U\hat{\mathfrak{g}}_\gamma / J_{\gamma,\lambda,k}, \quad (28)$$

where $J_{\gamma,\lambda,k}$ is the left ideal generated by $\mathfrak{n}_{\gamma,+}$, $H - \lambda(H)$ for $H \in \mathfrak{h}_\gamma$ and $\hat{k} - k$. The cyclic vector $1 \bmod J_{\gamma,\lambda,k}$ is denoted by 1_λ . Equivalently M_λ is defined as the $\hat{\mathfrak{g}}_\gamma$ -module induced from the one-dimensional $\hat{\mathfrak{b}}_\gamma$ -module:

$$M_\lambda := \text{Ind}_{\hat{\mathfrak{b}}_\gamma}^{\hat{\mathfrak{g}}_\gamma} \mathbb{C}1_\lambda, \quad (29)$$

where 1_λ satisfies $\mathfrak{n}_{\gamma,+}1_\lambda = 0$, $H1_\lambda = \lambda(H)1_\lambda$ and $\hat{k}1_\lambda = k1_\lambda$. (See Ref. [3], (1.3).)

(ii) The $\hat{\mathfrak{g}}_\gamma$ -module

$$M(0) := U\hat{\mathfrak{g}}_\gamma / J(0), \quad (30)$$

where $J(0) = U\hat{\mathfrak{g}}_\gamma \mathfrak{n}_{\gamma,+} + U\hat{\mathfrak{g}}_\gamma (\hat{k} - k)$, is called the *universal Verma* module of level k . [11] The cyclic vector $1 \bmod J(0)$ is denoted by $\mathbf{1}$. Equivalently $M(0)$ is defined as the $\hat{\mathfrak{g}}_\gamma$ -module induced from the one-dimensional $(\mathfrak{n}_{\gamma,+} \oplus \mathbb{C}\hat{k})$ -module:

$$M(0) := \text{Ind}_{\mathfrak{n}_{\gamma,+} \oplus \mathbb{C}\hat{k}}^{\hat{\mathfrak{g}}} \mathbb{C}\mathbf{1}, \quad (31)$$

where $\mathbf{1}$ satisfies $\mathfrak{n}_{\gamma,+}\mathbf{1} = 0$ and $\hat{k}\mathbf{1} = k\mathbf{1}$.

Since $[\mathfrak{h}_\gamma, J_{\gamma,\lambda,k}] \subset J_{\gamma,\lambda,k}$ and $[\mathfrak{h}_\gamma, J(0)] \subset J(0)$, we can define right $\mathfrak{h}_\gamma$ -actions on M_λ and $M(0)$ by

$$[x] \mapsto [xH], \quad (32)$$

for $H \in \mathfrak{h}_\gamma$ and $x \in U\hat{\mathfrak{g}}_\gamma$, where $[x]$ denotes the equivalence class of x modulo $J_{\gamma,\lambda,k}$ or $J(0)$.

The general factorisation theorem is as follows. Hereafter we always assume that $\{Q_1, \dots, Q_L\}$ do not contain neither 0 nor ∞ . Hence $D = Q_1 + \dots + Q_L$ is a divisor of E_{trig} as well as that of E_{orb} .

Theorem 4.2. *Let M_i be a smooth $\hat{\mathfrak{g}}^{Q_i}$ -module of level k for $i = 1, \dots, L$ and M be the tensor product $M_1 \otimes \dots \otimes M_L$ endowed with the $\hat{\mathfrak{g}}^D$ -module structure. Then,*

$$\begin{aligned} \text{CC}_{\text{trig}}(M) &\cong \text{CC}_{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty) / \text{CC}_{\text{orb}} \cdot \mathfrak{h}, \\ \text{CC}_{\text{orb}} \cdot \mathfrak{h} &:= \text{CC}_{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty) \rho_{1,\beta}(\mathfrak{h}). \end{aligned} \quad (33)$$

Here $M(0)_0$ and $M(0)_\infty$ are the universal Verma modules defined above for $\hat{\mathfrak{g}}_\gamma = \hat{\mathfrak{g}}^{(0)}$ and for $\hat{\mathfrak{g}}_\gamma = \hat{\mathfrak{g}}^{(\infty)}$ respectively and $\rho_{1,\beta}$ is the right action of $\mathfrak{h} = \mathfrak{h}^{(0)} = \mathfrak{h}^{(\infty)}$ on $M(0)_0 \otimes M \otimes M(0)_\infty$ defined by

$$(v_0 \otimes v \otimes v_\infty) \rho_{1,\beta}(H) := v_0 H \otimes v \otimes v_\infty + v_0 \otimes v \otimes v_\infty \text{Ad}(\beta)H \quad (34)$$

for $H \in \mathfrak{h}$, $v_0 \in M(0)_0$, $v \in M$ and $v_\infty \in M(0)_\infty$. Note that the left action of $\mathfrak{g}_{\text{out}}^{\text{orb}}$ and the right action of $\mathfrak{h}$ on $M(0)_0 \otimes M \otimes M(0)_\infty$ commute with each other and thus $\rho_{1,\beta}$ descends to the action on CC_{orb} .

This statement is rigorous but not very illuminating, so we rephrase it in a little bit incorrect but more impressive way. By virtue of the Poincaré-Birkhoff-Witt theorem, M_λ and $M(0)$ are isomorphic to $U\mathfrak{n}_{\gamma,-}$ and $U\mathfrak{n}_{\gamma,-} \otimes U\mathfrak{h}_\gamma$ respectively as linear spaces. Being commutative, the universal enveloping algebra $U\mathfrak{h}_\gamma$ is a polynomial ring over the dual space $\mathfrak{h}_\gamma^* = \text{Hom}_\mathbb{C}(\mathfrak{h}_\gamma, \mathbb{C})$. Hence $M(0)$ is spanned by $x \otimes f(\lambda)$, where $x \in U\mathfrak{g}_{\gamma,-}$ and $f(\lambda)$ is a polynomial in $\lambda \in \mathfrak{h}_\gamma^*$. Evaluating $x \otimes f(\lambda)$ at $\lambda \in \mathfrak{h}_\gamma^*$, we obtain an element of M_λ . In this sense, $M(0)$ may be approximately regarded as a direct integral:

$$M(0) \text{ “=” } \int_{\lambda \in \mathfrak{h}_\gamma^*} M_\lambda d\lambda. \quad (35)$$

(This is not correct since the direct integral should be spanned by L^2 -functions instead of polynomials.) Using this (approximate) identification, we can rewrite $\text{CC}_{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty)$ as

$$\text{CC}_{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty) = \iint_{\mathfrak{h}^* \times \mathfrak{h}^*} \text{CC}_{\text{orb}}(M_\lambda \otimes M \otimes M_\mu) d\lambda d\mu. \quad (36)$$

The statement of Theorem 4.2 becomes

$$\text{CC}_{\text{trig}}(M) \cong \int_{\mathfrak{h}} \text{CC}_{\text{orb}}(M_\lambda \otimes M \otimes M_{\beta^*(\lambda)}) d\lambda, \quad (37)$$

where $\beta^*(\lambda) = -\lambda \circ \text{Ad}(\beta^{-1})$. The proof of (37) (from Theorem 4.2) is the same as that of (72).

The isomorphism (37) is an analogue of the factorisation theorem of ordinary WZW models.[5, 12, 13, 9] In fact, under certain finiteness conditions for M_i 's, almost all $\text{CC}_{\text{orb}}(M_\lambda \otimes M \otimes M_{\beta^*(\lambda)})$ vanish and the direct integral is replaced by finite direct sum as we will prove in §5.

Proof. The idea of the proof of Theorem 4.2 is essentially the same as that of Ref. [5]. See also Ref. [12, 13]. We make use of techniques in Ref. [9].

To begin with, observe that there is an exact sequence,

$$0 \rightarrow \mathfrak{g}_{\text{out}}^{\text{trig}} M / \mathfrak{g}_{\text{out}}^0 M \rightarrow M / \mathfrak{g}_{\text{out}}^0 M \rightarrow \text{CC}_{\text{trig}}(M) \rightarrow 0, \quad (38)$$

where

$$\begin{aligned} \mathfrak{g}_{\text{out}}^0 &= \{f(t) : \text{meromorphic section of } \mathfrak{g}^{\text{orb}} \text{ with poles in } D, f(0) = f(\infty) = 0\} \\ &\subset \mathfrak{g}_{\text{out}}^{\text{trig}}. \end{aligned} \quad (39)$$

Using this exact sequence, we prove the theorem in two steps. In the first step, we show that a linear map $\varphi(v \bmod \mathfrak{g}_{\text{out}}^0 M) = [\mathbf{1}_0 \otimes v \otimes \mathbf{1}_\infty] \in \text{CC}_{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty)$ induces an isomorphism

$$\varphi : M/\mathfrak{g}_{\text{out}}^0 M \xrightarrow{\sim} \text{CC}_{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty). \quad (40)$$

In the second step we show that

$$\varphi(\mathfrak{g}_{\text{out}}^{\text{trig}} M/\mathfrak{g}_{\text{out}}^0 M) \cong \text{CC}_{\text{orb}} \cdot \mathfrak{h}, \quad (41)$$

(cf. (33) for the definition of $\text{CC}_{\text{orb}} \cdot \mathfrak{h}$) or, equivalently,

$$\mathbf{1}_0 \otimes \mathfrak{g}_{\text{out}}^{\text{trig}} M \otimes \mathbf{1}_\infty \equiv (M(0)_0 \otimes M \otimes M(0)_\infty) \rho_{1,\beta}(\mathfrak{h}) \bmod \mathfrak{g}_{\text{out}}^{\text{orb}}. \quad (42)$$

Hereafter $\bmod \mathfrak{g}_{\text{out}}^{\text{orb}}$ denotes $\bmod \mathfrak{g}_{\text{out}}^{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty)$. Combining these two statements with the exact sequence (38), we obtain the proof of the theorem.

To prove that φ in (40) is an isomorphism, we use a lemma by Kazhdan and Lusztig.[14]

Lemma 4.3. *Let $\mathfrak{a}$ be a Lie algebra over $\mathbb{C}$ and $\mathfrak{a}_1, \mathfrak{a}_2$ be its Lie subalgebra such that $\mathfrak{a} = \mathfrak{a}_1 + \mathfrak{a}_2$. Assume that an $\mathfrak{a}_2$ -module V is given and put $W := \text{Ind}_{\mathfrak{a}_2}^{\mathfrak{a}} V$. Then the canonical homomorphism $V \ni v \mapsto v \in W$ induces an isomorphism*

$$V/(\mathfrak{a}_1 \cap \mathfrak{a}_2)V \xrightarrow{\sim} W/\mathfrak{a}_1 W. \quad (43)$$

See §A.7 of Ref. [14] for the proof.

We apply this lemma to

$$\mathfrak{a}_1 = \mathfrak{g}_{\text{out}}^{\text{orb}}, \quad \mathfrak{a}_2 = \mathfrak{n}_+^{(0)} \oplus \mathfrak{g}^D \oplus \mathfrak{n}_+^{(\infty)} \oplus \mathbb{C}\hat{k}, \quad V = \mathbb{C}\mathbf{1}_0 \otimes M \otimes \mathbb{C}\mathbf{1}_\infty, \quad (44)$$

where $\mathbf{1}_0$ and $\mathbf{1}_\infty$ are the canonical cyclic vectors of $M(0)_0$ and $M(0)_\infty$. It is easy to see from (26) or (27) that the Lie algebra $\mathfrak{a} = \mathfrak{a}_1 + \mathfrak{a}_2$ is equal to $\mathfrak{g}^{(0)} \oplus \mathfrak{g}^D \oplus \mathfrak{g}^{(\infty)} \oplus \mathbb{C}\hat{k}$. Since $\mathfrak{a}_1 \cap \mathfrak{a}_2 = \mathfrak{g}_{\text{out}}^0$ acts trivially on the first and the last factor of $M(0)_0 \otimes M \otimes M(0)_\infty$, we have

$$V/(\mathfrak{a}_1 \cap \mathfrak{a}_2)V \cong \mathbb{C}\mathbf{1}_0 \otimes M \otimes \mathbb{C}\mathbf{1}_\infty / \mathfrak{g}_{\text{out}}^0 \mathbb{C}(\mathbf{1}_0 \otimes M \otimes \mathbb{C}\mathbf{1}_\infty) = M/\mathfrak{g}_{\text{out}}^0 M. \quad (45)$$

On the other hand,

$$W = \text{Ind}_{\mathfrak{n}_+^{(0)} \oplus \mathfrak{g}^D \oplus \mathfrak{n}_+^{(\infty)} \oplus \mathbb{C}\hat{k}}^{\mathfrak{g}^{(0)} \oplus \mathfrak{g}^D \oplus \mathfrak{g}^{(\infty)} \oplus \mathbb{C}\hat{k}} \mathbb{C}\mathbf{1}_0 \otimes M \otimes \mathbb{C}\mathbf{1}_\infty = M(0)_0 \otimes M \otimes M(0)_\infty. \quad (46)$$

Hence $W/\mathfrak{a}_1 W = M(0)_0 \otimes M \otimes M(0)_\infty / \mathfrak{g}_{\text{out}}^{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty)$, which proves (40) by Lemma 4.3.

Next we prove (42). Namely, we prove the following two claims:

$$\mathbf{1}_0 \otimes \mathfrak{g}_{\text{out}}^{\text{trig}} M \otimes \mathbf{1}_\infty \subset (M(0)_0 \otimes M \otimes M(0)_\infty) \rho_{1,\beta}(\mathfrak{h}) + \mathfrak{g}_{\text{out}}^{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty). \quad (47)$$

$$\mathbf{1}_0 \otimes \mathfrak{g}_{\text{out}}^{\text{trig}} M \otimes \mathbf{1}_\infty + \mathfrak{g}_{\text{out}}^{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty) \supset (M(0)_0 \otimes M \otimes M(0)_\infty) \rho_{1,\beta}(\mathfrak{h}). \quad (48)$$

To prove (47), assume $f(t) \in \mathfrak{g}_{\text{out}}^{\text{trig}}$ and $v \in M$. We show that $\mathbf{1}_0 \otimes f(t)v \otimes \mathbf{1}_\infty$ belongs to $(M(0)_0 \otimes M \otimes M(0)_\infty) \rho_{1,\beta}(\mathfrak{h}) \pmod{\mathfrak{g}_{\text{out}}^{\text{orb}}}$. Since $f(t) \in \mathfrak{g}_{\text{out}}^{\text{trig}} \subset \mathfrak{g}_{\text{out}}^{\text{orb}}$, we have

$$\mathbf{1}_0 \otimes f(t)v \otimes \mathbf{1}_\infty \equiv -f(0)\mathbf{1}_0 \otimes v \otimes \mathbf{1}_\infty - \mathbf{1}_0 \otimes v \otimes f(\infty)\mathbf{1}_\infty \pmod{\mathfrak{g}_{\text{out}}^{\text{orb}}}. \quad (49)$$

The conditions (17) and (18) imply that $f(\infty) = \text{Ad } \beta(f(0)) \in \mathfrak{h}$. Hence the right hand side of (49) belongs to $(M(0)_0 \otimes M \otimes M(0)_\infty) \rho_{1,\beta}(\mathfrak{h})$ which proves (47).

Conversely, assume $v_0 \in M(0)_0$, $v \in M$, $v_\infty \in M(0)_\infty$ and $H \in \mathfrak{h}$. We show that $(v_0 \otimes v \otimes v_\infty) \rho_{1,\beta}(H)$ belongs to $\mathbf{1}_0 \otimes \mathfrak{g}_{\text{out}}^{\text{trig}} M \otimes \mathbf{1}_\infty \pmod{\mathfrak{g}_{\text{out}}^{\text{orb}}}$ to prove (48). Thanks to the fact we proved in the first step of this proof of Theorem 4.2, there exists $\tilde{v} \in M$ such that

$$v_0 \otimes v \otimes v_\infty \equiv \mathbf{1}_0 \otimes \tilde{v} \otimes \mathbf{1}_\infty \pmod{\mathfrak{g}_{\text{out}}^{\text{orb}}}. \quad (50)$$

Hence we have

$$\begin{aligned} (v_0 \otimes v \otimes v_\infty) \rho_{1,\beta}(H) &\equiv (\mathbf{1}_0 \otimes \tilde{v} \otimes \mathbf{1}_\infty) \rho_{1,\beta}(H) \\ &= H \otimes \tilde{v} \otimes \mathbf{1}_\infty + \mathbf{1}_0 \otimes \tilde{v} \otimes \text{Ad } \beta(H) \\ &\equiv -\mathbf{1}_0 \otimes h(t)\tilde{v} \otimes \mathbf{1}_\infty \end{aligned} \quad (51)$$

modulo $\mathfrak{g}_{\text{out}}^{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty)$. Here $h(t)$ is an element of $\mathfrak{g}_{\text{out}}^{\text{trig}}$ satisfying $h(0) = H + (\text{regular function at } t = 0)$ and $h(\infty) = \text{Ad } \beta(H) + (\text{regular function at } t = \infty)$. When H is $J_{0,b}$ (cf. (7)), such $h(t)$ can be written down explicitly:

$$h_b(t) := \frac{\varepsilon^b t^N - t_i^N}{t^N - t_i^N} J_{0,b}, \quad (52)$$

where $t_i = \exp(2\pi i z_i / N)$ is the coordinate of Q_i for a certain i ($1 \leq i \leq N$). For general H , $h(t)$ can be constructed as a linear combination of $h_b(t)$. The last expression in (51) is an element of $\mathbf{1}_0 \otimes \mathfrak{g}_{\text{out}}^{\text{trig}} M \otimes \mathbf{1}_\infty$, which proves (48). Thus (42) as well as Theorem 4.2 is proved. $\square$

5 Factorisation (2)

The statement of the previous section can be refined for special modules. In fact, the direct integral in (37) is replaced by finite direct sum when we impose following finiteness conditions on M_i 's: for each $i = 1, \dots, L$ there exists a finite dimensional $\mathfrak{g}$ -module V_i which generates M_i over $\hat{\mathfrak{g}}^{Q_i}$ and $\hat{\mathfrak{g}}_+^{Q_i} V_i \subset V_i$. Throughout this section we assume this condition, which automatically implies that M_i 's are smooth.

Let us denote $V = V_1 \otimes \dots \otimes V_L$ and the set of $\mathfrak{g}$ -weights of V by $\text{wt}(V)$. For $\lambda \in \mathfrak{h}^*$, we define

$$\begin{aligned}\tilde{\lambda} &:= -\lambda \circ (1 - \text{Ad } \beta^{-1})^{-1} = \lambda \circ (1 - \text{Ad } \beta)^{-1} \circ \text{Ad } \beta, \\ \tilde{\lambda}' &:= -\lambda \circ (1 - \text{Ad } \beta)^{-1}.\end{aligned}\tag{53}$$

They are well-defined since $1 - \text{Ad } \beta^{\pm 1}$ is invertible on $\mathfrak{h} = \bigoplus_{b=1}^{N-1} \mathbb{C} J_{0,b}$.

Theorem 5.1. *Under the above assumption on M_i , $\text{CC}_{\text{trig}}(M)$ is finite dimensional and there is a linear isomorphism*

$$\text{CC}_{\text{trig}}(M) \cong \bigoplus_{\lambda \in \text{wt}(V)} \text{CC}_{\text{orb}}(M_{\tilde{\lambda}} \otimes M \otimes M_{\tilde{\lambda}'}),\tag{54}$$

where $M_{\tilde{\lambda}}$ and $M_{\tilde{\lambda}'}$ are inserted at 0 and ∞ of E_{orb} respectively.

Proof. For simplicity of notation, we denote $J_{0,b}$ by H_b ($b = 1, \dots, N-1$). As mentioned before, they form a basis of $\mathfrak{h}$. We also introduce ideals $I_{\mathfrak{h}}$ and $I'_{\mathfrak{h}}$ of the polynomial ring $U\mathfrak{h}$ as follows:

$$\begin{aligned}I_{\mathfrak{h}} &= \bigcap_{\lambda \in \text{wt}(V)} I_{\lambda}, & I'_{\mathfrak{h}} &= \bigcap_{\lambda \in \text{wt}(V)} I'_{\lambda}, \\ I_{\lambda} &:= \text{ideal of } U\mathfrak{h} \text{ generated by } H_b - \tilde{\lambda}(H_b) (b = 1, \dots, N-1). \\ I'_{\lambda} &:= \text{ideal of } U\mathfrak{h} \text{ generated by } H_b - \tilde{\lambda}'(H_b) (b = 1, \dots, N-1).\end{aligned}\tag{55}$$

The main part of the proof of the theorem is to show the following inclusion:

$$I \otimes M \otimes M(0)_{\infty} + M(0)_0 \otimes M \otimes I' \subset K,\tag{56}$$

where

$$I = M(0)_0 I_{\mathfrak{h}}, \quad I' = M(0)_{\infty} I'_{\mathfrak{h}},\tag{57}$$

and K is the kernel of the canonical surjection $M(0)_0 \otimes M \otimes M(0)_\infty \rightarrow \text{CC}_{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty) / \text{CC}_{\text{orb}} \cdot \mathfrak{h}$, namely,

$$K := \mathfrak{g}_{\text{out}}^{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty) + (M(0)_0 \otimes M \otimes M(0)_\infty) \rho_{1,\beta}(\mathfrak{h}). \quad (58)$$

When the inclusion (56) is proved, we shall see that we may replace $\text{CC}_{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty)$ in (33) by $\text{CC}_{\text{orb}}(\bar{M}(0)_0 \otimes M \otimes \bar{M}(0)_\infty)$, where $\bar{M}(0)_0 = M(0)_0 / I$ and $\bar{M}(0)_\infty = M(0)_\infty / I'$ which are finite direct sums of Verma modules. The space of conformal coinvariants $\text{CC}_{\text{orb}}(\bar{M}(0)_0 \otimes M \otimes \bar{M}(0)_\infty)$ decomposes into finite direct sum of $\text{CC}_{\text{orb}}(M_{\bar{\lambda}} \otimes M \otimes M_{\bar{\lambda}'})$ and, factoring by $\text{CC}_{\text{orb}} \cdot \mathfrak{h}$, we shall obtain (54). The proof of the first statement, $\dim \text{CC}_{\text{trig}}(M) < \infty$, is given after Lemma 5.2.

Let us show that

$$I \otimes M \otimes M(0)_\infty \subset K, \quad (59)$$

namely, for $v_0 \otimes v \otimes v_\infty \in M(0)_0 \otimes M \otimes M(0)_\infty$ and $p(H) \in I_{\mathfrak{h}}$, $v_0 p(H) \otimes v \otimes v_\infty$ belongs to K . Thanks to the existence of $\tilde{v}$ satisfying (50), we may assume $v_0 = \mathbf{1}_0$ and $v_\infty = \mathbf{1}_\infty$, because $\mathfrak{g}_{\text{out}}^{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty)(p(H) \otimes \mathbf{1} \otimes \mathbf{1}) \subset K$. Hence the problem reduces to show

$$p(H) \otimes v \otimes \mathbf{1}_\infty \in K. \quad (60)$$

By the following lemma, we have only to show (60) for $v \in V \subset M$.

Lemma 5.2. (i) $\hat{\mathfrak{g}}^D = \mathfrak{g}_{\text{out}}^{\text{trig}} \oplus \mathfrak{g}_+^D \oplus \mathbb{C}\hat{k}$. (ii) $M = V + \mathfrak{g}_{\text{out}}^{\text{trig}} M$.

Proof. (i) The space $\mathfrak{g}^D$ is spanned by $\mathfrak{g}_+^D$ and the elements of the form $0 \oplus \cdots \oplus J_{a,b} \otimes \xi_i^{-n} \oplus \cdots \oplus 0$, where $(a, b) \in \{0, 1, \dots, N-1\}^2 \setminus \{(0, 0)\}$, $n \in \mathbb{Z}_{>0}$, ξ_i is the local coordinate at $Q_i \in D$ and $J_{a,b} \otimes \xi_i^{-n}$ sits at the i -th component. The latter elements can be replaced with the following which belong to $\mathfrak{g}_{\text{out}}^{\text{trig}}$: if $n = 1$,

$$J_{a,b}(t) := \begin{cases} J_{a,b} t^a (t^N - t_i^N)^{-1} & (a \neq 0), \\ J_{0,b} (\varepsilon^b t^N - t_i^N) (t^N - t_i^N)^{-1} & (a = 0). \end{cases} \quad (61)$$

and if $n > 1$, $(t d/dt)^{n-1} J_{a,b}(t)$. Thus we have $\mathfrak{g}_{\text{out}}^{\text{trig}} + \mathfrak{g}_+^D = \mathfrak{g}^D$. It is an exercise of complex analysis to show that $\mathfrak{g}_{\text{out}}^{\text{trig}} \cap \mathfrak{g}_+^D = \{0\}$.

(ii) By the assumption $M_i = U \hat{\mathfrak{g}}^{Q_i} V_i$, $\hat{\mathfrak{g}}_+^{Q_i} V_i \subset V_i$ and the Poincaré-Birkhoff-Witt theorem, we have $M = U \hat{\mathfrak{g}}^D V = U \mathfrak{g}_{\text{out}}^{\text{trig}} V = V + \mathfrak{g}_{\text{out}}^{\text{trig}} M$. $\square$

By this lemma, every element $v \in M$ is decomposed as $v = v_f + v_{\text{out}}$, $v_f \in V$, $v_{\text{out}} \in \mathfrak{g}_{\text{out}}^{\text{trig}} M$. The finite-dimensionality of $\text{CC}_{\text{trig}}(M)$ immediately follows from this expression. Note that $\mathfrak{g}_{\text{out}}^{\text{trig}}$ acts trivially on $\mathbf{1}_0$ and $\mathbf{1}_\infty$. Hence $\mathbf{1}_0 \otimes v_{\text{out}} \otimes \mathbf{1}_\infty$ belongs to $\mathfrak{g}_{\text{out}}^{\text{orb}}(M(0)_0 \otimes M \otimes M(0)_\infty) \subset K$. K becomes a right $U\mathfrak{h}$ -module by the action $v_0 \otimes v \otimes v_\infty \mapsto v_0 H \otimes v \otimes v_\infty$ of $H \in \mathfrak{h}$. Therefore $p(H) \otimes v_{\text{out}} \otimes \mathbf{1}_\infty$ belongs to K , which allows us to assume that $v \in V$ in (60).

We further reduces the problem and show (60) for a weight vector in V , $v_\lambda \in V_\lambda$. K being a right $U\mathfrak{h}$ -module, as we mentioned above, it is enough to show that

$$(H_b - \tilde{\lambda}(H_b)) \otimes v_\lambda \otimes \mathbf{1}_\infty \in K, \quad (62)$$

for $b = 1, \dots, N-1$. Since the constant function with value H_b belongs to $\mathfrak{g}_{\text{out}}^{\text{orb}}$, we have

$$\begin{aligned} H_b \otimes v_\lambda \otimes \mathbf{1}_\infty &\equiv -\mathbf{1}_0 \otimes H_b v_\lambda \otimes \mathbf{1}_\infty - \mathbf{1}_0 \otimes v_\lambda \otimes H_b \\ &= -\mathbf{1}_0 \otimes \lambda(H_b) v_\lambda \otimes \mathbf{1}_\infty - \mathbf{1}_0 \otimes v_\lambda \otimes \text{Ad } \beta(\text{Ad } \beta^{-1}(H_b)) \\ &\equiv -\lambda(H_b) \mathbf{1}_0 \otimes v_\lambda \otimes \mathbf{1}_\infty + \text{Ad } \beta^{-1}(H_b) \otimes v_\lambda \otimes \mathbf{1}_\infty \\ &= -\lambda(H_b) \mathbf{1}_0 \otimes v_\lambda \otimes \mathbf{1}_\infty + \varepsilon^{-b} H_b \otimes v_\lambda \otimes \mathbf{1}_\infty \end{aligned} \quad (63)$$

modulo K , which proves (62) and consequently (59). Similarly we can prove that $M(0)_0 \otimes M \otimes I' \subset K$ and the inclusion (56) is proved.

Now we have the following commutative diagram with exact rows.

$$\begin{array}{ccccccc} & 0 & & 0 & & & \\ & \downarrow & & \downarrow & & & \\ 0 & \rightarrow \tilde{I} & = & \tilde{I} & \rightarrow & 0 & \\ & \downarrow & & \downarrow & & \downarrow & \\ 0 & \rightarrow K & \rightarrow & M(0)_0 \otimes M \otimes M(0)_\infty & \rightarrow & \text{CC}_{\text{trig}}(M) & \rightarrow 0 \\ & \downarrow & & \downarrow & & \downarrow & \\ 0 & \rightarrow \bar{K} & \rightarrow & \bar{M}(0)_0 \otimes M \otimes \bar{M}(0)_\infty & \rightarrow & \bar{M}(0)_0 \otimes M \otimes \bar{M}(0)_\infty / \bar{K} & \rightarrow 0 \\ & \downarrow & & \downarrow & & \downarrow & \\ & 0 & & 0 & & 0 & \end{array} \quad (64)$$

Here $\tilde{I} := I \otimes M \otimes M(0)_\infty + M(0)_0 \otimes M \otimes I'$ and $\bar{K} := K / \tilde{I}$. (The exactness of the middle row is due to Theorem 4.2.) Applying the nine lemma in the

homological algebra to this diagram (or chasing the diagram), we have

$$\mathrm{CC}_{\mathrm{trig}}(M) \cong \bar{M}(0)_0 \otimes M \otimes \bar{M}(0)_\infty / \bar{K}. \quad (65)$$

It is easy to see that

$$\bar{K} = \mathfrak{g}_{\mathrm{out}}^{\mathrm{orb}}(\bar{M}(0)_0 \otimes M \otimes \bar{M}(0)_\infty) + (\bar{M}(0)_0 \otimes M \otimes \bar{M}(0)_\infty)\rho_{1,\beta}(\mathfrak{h}), \quad (66)$$

because the left $\hat{\mathfrak{g}}^D$ -action and the right $\mathfrak{h}$ -action on $M(0)_0 \otimes M \otimes M(0)_\infty$ commute and the right $\mathfrak{h}$ -action is commutative. By (65) and (66) we have

$$\begin{aligned} \mathrm{CC}_{\mathrm{trig}}(M) &\cong \mathrm{CC}_{\mathrm{orb}}(\bar{M}(0)_0 \otimes M \otimes \bar{M}(0)_\infty) / \overline{\mathrm{CC}_{\mathrm{orb}}} \cdot \rho_{1,\beta}(\mathfrak{h}), \\ \overline{\mathrm{CC}_{\mathrm{orb}}} \cdot \rho_{1,\beta}(\mathfrak{h}) &:= \mathrm{CC}_{\mathrm{orb}}(\bar{M}(0)_0 \otimes M \otimes \bar{M}(0)_\infty)\rho_{1,\beta}(\mathfrak{h}). \end{aligned} \quad (67)$$

Lemma 5.3.

$$\bar{M}(0)_0 \cong \bigoplus_{\lambda \in \mathrm{wt}(V)} M_{\tilde{\lambda}}, \quad \bar{M}(0)_\infty \cong \bigoplus_{\lambda \in \mathrm{wt}(V)} M_{\tilde{\lambda}'}. \quad (68)$$

(cf. (53).)

Proof. For two distinct weights λ and μ in $\mathrm{wt}(V)$, we have $I_\lambda + I_\mu = U\mathfrak{h}$, since $\{H_b\}_{b=1,\dots,N-1}$ is a basis of $\mathfrak{h}$ and the map $\lambda \mapsto \tilde{\lambda}$ is injective. (See (55).) Hence the “Chinese remainder theorem” implies

$$U\mathfrak{h}/I_{\mathfrak{h}} \cong \bigoplus_{\lambda \in \mathrm{wt}(V)} U\mathfrak{h}/I_\lambda. \quad (69)$$

Regarding both hand sides as $(\mathfrak{g}_+^{(0)} \oplus \mathfrak{h} \oplus \mathbb{C}\hat{k})$ -modules, on which $\mathfrak{g}_+^{(0)}$ acts trivially and $\hat{k}$ acts as k , we induce them up to $\hat{\mathfrak{g}}^{(0)}$ -modules and we obtain the first isomorphism in (68). The second isomorphism in (68) is proved in the same manner. $\square$

Due to Lemma 5.3 we may rewrite (67) as

$$\begin{aligned} \mathrm{CC}_{\mathrm{trig}}(M) &\cong \bigoplus_{\lambda, \mu \in \mathrm{wt}(V)} \mathrm{CC}_{\mathrm{orb}}(M_{\tilde{\lambda}} \otimes M \otimes M_{\tilde{\mu}'}) / \mathrm{CC}_{\mathrm{orb}, \lambda, \mu} \cdot \rho_{1,\beta}(\mathfrak{h}), \\ \mathrm{CC}_{\mathrm{orb}, \lambda, \mu} \cdot \rho_{1,\beta}(\mathfrak{h}) &:= \mathrm{CC}_{\mathrm{orb}}(M_{\tilde{\lambda}} \otimes M \otimes M_{\tilde{\mu}'})\rho_{1,\beta}(\mathfrak{h}). \end{aligned} \quad (70)$$

We show that the component $\mathrm{CC}_{\mathrm{orb}}(M_{\tilde{\lambda}} \otimes M \otimes M_{\tilde{\mu}'}) / \mathrm{CC}_{\mathrm{orb}, \lambda, \mu} \cdot \rho_{1,\beta}(\mathfrak{h})$ of the above decomposition vanishes unless $\lambda = \mu$. Let us take $v_0 \otimes v \otimes v_\infty \in$

$M_{\tilde{\lambda}} \otimes M \otimes M_{\tilde{\mu}'}$. For any $H \in \mathfrak{h}$, we have $v_0 H = \tilde{\lambda}(H)v_0$ and $v_\infty H = \tilde{\mu}'(H)v_\infty$, which lead to

$$\begin{aligned} (v_0 \otimes v \otimes v_\infty) \rho_{1,\beta}(H) &= v_0 H \otimes v \otimes v_\infty + v_0 \otimes v \otimes v_\infty \text{Ad}(\beta)H \\ &= (\lambda - \mu)(1 - \text{Ad} \beta)^{-1}(H)(v_0 \otimes v \otimes v_\infty), \end{aligned} \quad (71)$$

due to (53). Hence if $\lambda - \mu \neq 0$, there exists $H \in \mathfrak{h}$ such that $(\lambda - \mu)(1 - \text{Ad} \beta)^{-1}(H) \neq 0$, which means $v_0 \otimes v \otimes v_\infty \in \text{CC}_{\text{orb},\lambda,\mu} \cdot \rho_{1,\beta}(\mathfrak{h})$. Therefore

$$\begin{aligned} \text{CC}_{\text{orb}}(M_{\tilde{\lambda}} \otimes M \otimes M_{\tilde{\mu}'}) / \text{CC}_{\text{orb},\lambda,\mu} \cdot \rho_{1,\beta}(\mathfrak{h}) \\ = \begin{cases} \text{CC}_{\text{orb}}(M_{\tilde{\lambda}} \otimes M \otimes M_{\tilde{\lambda}'}) & (\lambda = \mu), \\ 0 & (\lambda \neq \mu). \end{cases} \end{aligned} \quad (72)$$

This completes the proof of Theorem 5.1. $\square$

6 Example (case of Weyl modules)

Let us directly compute the space of conformal coinvariants for Weyl modules as a simple example and see how the factorisation theorem Theorem 5.1 decompose CC_{trig} into CC_{orb} 's.

Let V_i be a finite-dimensional $\mathfrak{g}$ -module and regard them as $(\mathfrak{g}_+^{Q_i} \oplus \mathbb{C}\hat{k})$ -module of level k as before. The *Weyl module* is the $\hat{\mathfrak{g}}^{Q_i}$ -module induced up from V_i :

$$M(V_i) := \text{Ind}_{\mathfrak{g}_+^{Q_i}}^{\mathfrak{g}_+^{Q_i} \oplus \mathbb{C}\hat{k}} V_i. \quad (73)$$

The tensor product $M(V) = M(V_1) \otimes \cdots \otimes M(V_L)$ is naturally induced up from $V = V_1 \otimes \cdots \otimes V_L$ and, by virtue of Lemma 5.2 (i), we have

$$M(V) = \text{Ind}_{\mathfrak{g}_+^D}^{\mathfrak{g}_+^D \oplus \mathbb{C}\hat{k}} V = U \mathfrak{g}_{\text{out}}^{\text{trig}} \otimes_{\mathbb{C}} V = V \oplus \mathfrak{g}_{\text{out}}^{\text{trig}} U \mathfrak{g}_{\text{out}}^{\text{trig}} \otimes_{\mathbb{C}} V = V \oplus \mathfrak{g}_{\text{out}}^{\text{trig}} M(V). \quad (74)$$

Hence the space of conformal coinvariants is isomorphic to V itself:

$$\text{CC}_{\text{trig}}(M(V)) \cong V. \quad (75)$$

On the other hand, let us compute the component $\text{CC}_{\text{orb}}(M_{\tilde{\lambda}} \otimes M(V) \otimes M_{\tilde{\lambda}'})$ of the factorisation, (54). Put $\mathfrak{a}_1 = \mathfrak{g}_{\text{out}}^{\text{orb}}$, $\mathfrak{a}_2 = \mathfrak{b}^{(0)} \oplus \mathfrak{b}^D \oplus \mathfrak{b}^{(\infty)} \oplus \mathbb{C}\hat{k}$,

where $\mathfrak{b}^{(0)}$ and $\mathfrak{b}^{(\infty)}$ are subalgebras of $\hat{\mathfrak{g}}^{(0)}$ and $\hat{\mathfrak{g}}^{(\infty)}$ defined at the beginning of §4 and $\mathfrak{b}^D := \bigoplus_{i=1}^L \mathfrak{g}_+^{Q_i}$. Then

$$\mathfrak{a} := \mathfrak{a}_1 + \mathfrak{a}_2 = \hat{\mathfrak{g}}^{D+(0)+(\infty)}, \quad \mathfrak{a}_1 \cap \mathfrak{a}_2 = \mathfrak{h}. \quad (76)$$

We apply Lemma 4.3 to these algebras and the $\mathfrak{a}_2$ -module $V \cong \mathbb{C}1_{\tilde{\lambda}} \otimes V \otimes \mathbb{C}1_{\tilde{\lambda}'}$. The module W in the lemma is

$$\text{Ind}_{\mathfrak{a}_2}^{\mathfrak{a}} \mathbb{C}1_{\tilde{\lambda}} \otimes V \otimes \mathbb{C}1_{\tilde{\lambda}'} = M_{\tilde{\lambda}} \otimes M(V) \otimes M_{\tilde{\lambda}'}. \quad (77)$$

The space of coinvariants “ $V/\mathfrak{a}_1 \cap \mathfrak{a}_2 V$ ” in the lemma is

$$\mathbb{C}1_{\tilde{\lambda}} \otimes V \otimes \mathbb{C}1_{\tilde{\lambda}'} / \mathfrak{h}(\mathbb{C}1_{\tilde{\lambda}} \otimes V \otimes \mathbb{C}1_{\tilde{\lambda}'}). \quad (78)$$

Take a weight vector $v_{\mu} \in V$ of weight μ . The action of $H \in \mathfrak{h}$ on $1_{\tilde{\lambda}} \otimes v_{\mu} \otimes 1_{\tilde{\lambda}'}$ is equal to the multiplication of $\tilde{\lambda}(H) + \mu(H) + \tilde{\lambda}'(H) = (-\lambda + \mu)(H)$. (See (53).) Namely, if $\lambda = \mu$, $\mathfrak{h}(1_{\tilde{\lambda}} \otimes v_{\mu} \otimes 1_{\tilde{\lambda}'}) = 0$, while if $\lambda \neq \mu$, $\mathfrak{h}(\mathbb{C}1_{\tilde{\lambda}} \otimes V \otimes \mathbb{C}1_{\tilde{\lambda}'})$ contains $1_{\tilde{\lambda}} \otimes v_{\mu} \otimes 1_{\tilde{\lambda}'}$. Consequently,

$$\mathbb{C}1_{\tilde{\lambda}} \otimes V \otimes \mathbb{C}1_{\tilde{\lambda}'} / \mathfrak{h}(\mathbb{C}1_{\tilde{\lambda}} \otimes V \otimes \mathbb{C}1_{\tilde{\lambda}'}) \cong V_{\lambda}. \quad (79)$$

The coinvariants “ $W/\mathfrak{a}_1 W$ ” in Lemma 4.3 is

$$M_{\tilde{\lambda}} \otimes M(V) \otimes M_{\tilde{\lambda}'} / \mathfrak{g}_{\text{out}}^{\text{orb}} M_{\tilde{\lambda}} \otimes M(V) \otimes M_{\tilde{\lambda}'} = \text{CC}_{\text{orb}}(M_{\tilde{\lambda}} \otimes M(V) \otimes M_{\tilde{\lambda}'}). \quad (80)$$

because of (77). Thus we have proved

$$\text{CC}_{\text{orb}}(M_{\tilde{\lambda}} \otimes M(V) \otimes M_{\tilde{\lambda}'}) \cong V_{\lambda}. \quad (81)$$

Summarising, the factorisation (54) for the Weyl modules $M_i = M(V_i)$ is the same as the weight space decomposition of V , $V = \bigoplus_{\lambda \in \text{wt}(V)} V_{\lambda}$.

7 Summary and concluding remarks

We formulated the twisted WZW model on a degenerate elliptic curve E_{trig} and the orbifold E_{orb} , Definition 3.2. Theorem 4.2 asserts that the space of conformal coinvariants (and thus the space of conformal blocks) on E_{trig} factorises into that on E_{orb} . This factorisation theorem holds for any smooth modules, but the result can be refined for modules generated by $\mathfrak{g}$ -modules, Theorem 5.1. The factorisation for the case of Weyl modules is nothing more than the weight space decomposition of the tensor product of $\mathfrak{g}$ -module, (75), (81).

From these results arise several problems:

- In Ref. [5] the factorisation of the space of conformal coinvariants/blocks on singular curves together with the existence of a flat connection (the KZ connection) leads to the locally freeness of the sheaf of conformal coinvariants/blocks. One of our next tasks is to follow this line to analyse the structure of the KZ connection, which might help finding the integral representation of the solutions of the elliptic KZ equations.
- The weights $\tilde{\lambda}$ and $\tilde{\lambda}'$ in Theorem 5.1 look quite strange. For example, $\tilde{\lambda} = -\lambda/2$ when $\mathfrak{g} = sl_2(\mathbb{C})$ ($N = 2$) and $M_{\tilde{\lambda}}$ can hardly be integrable when M_{λ} is. But such weights are inevitable due to the charge conservation. The same kind of phenomenon is implicit but exists also in Ref. [3], where the solution of the elliptic KZ equations is constructed as a twisted trace of vertex operators. Representation theoretical meaning of $M_{\tilde{\lambda}}$ is in question.
- We computed the space of conformal coinvariants for the Weyl modules in §6. This case is important in the study of the elliptic Gaudin model.[4] The case for integrable modules instead of Weyl modules would also be important.

Acknowledgments

The author expresses his gratitude to Akihiro Tsuchiya who explained details of his works, Michio Jimbo, Tetsuji Miwa, Hiroyuki Ochiai, Kiyoshi Ohba, Nobuyoshi Takahashi and Tomohide Terasoma for discussion and comments. He also thanks Gen Kuroki for discussions during the previous collaborations. Christian Schweigert and Martin Halpern who informed the references for the orbifold WZW model are gratefully acknowledged.

The atmosphere and environment of Mathematical Sciences Research Institute (Berkeley, U.S.A.), Institute for Theoretical and Experimental Physics (Moscow, Russia) and the VIth International Workshop “Conformal Field Theory and Integrable Models” (Chernogolovka, Russia) were very important. The author thanks their hospitality.

This work is partly supported by the Grant-in-Aid for Scientific Research, Japan Society for the Promotion of Science.

References

- [1] R. J. Baxter, *Ann. Phys.* **76**, 1–24, 25–47, 48–71 (1973).
- [2] A. A. Belavin, V. G. Drinfeld, *Funkts. Anal. i ego Prilozh.* **16-3**, 1–29 (1982) (in Russian); *Funct. Anal. Appl.* **16**, 159–180 (1982) (English transl.)
- [3] P. I. Etingof, *Comm. Math. Phys.* **159**, 471–502 (1994).
- [4] G. Kuroki, T. Takebe, *Comm. Math. Phys.* **190**, 1–56 (1997).
- [5] A. Tsuchiya, K. Ueno, Y. Yamada, In: *Integrable systems in quantum field theory and statistical mechanics*, *Adv. Stud. Pure Math.* **19**, 459–566 (1989).
- [6] S. Hamidi, C. Vafa, *Nucl. Phys.* **B279** 465–513 (1987); L. Borisov, M. B. Halpern, C. Schweigert, *Internat. J. Modern Phys.* **A13** 125–168, (1998); M. B. Halpern, J. E. Wang, 97–162 (2001); O. Ganor, M. B. Halpern, C. Helfgott, N. A. Obers, *J. High Energy Phys.* **0212** 019, (2002); M. B. Halpern, C. Helfgott, Twisted Einstein tensors and orbifold geometry, preprint [hep-th/0212275](#), to appear in *Int. J. Modern Phys. A*; J. de Boer, M. B. Halpern, N. A. Obers, *J. High Energy Phys.* **0110** 011, (2001).
- [7] E. Frenkel, D. Ben-Zvi, *Vertex algebras and algebraic curves* (American Mathematical Society, Providence, 2001) Mathematical Surveys and Monographs, **88**.
- [8] E. Frenkel, M. Szczesny, Twisted Modules over Vertex Algebras on Algebraic Curves, preprint [math.AG/0112211](#).
- [9] K. Nagatomo, A. Tsuchiya, Conformal field theories associated to regular chiral vertex operator algebras I: theories over the projective line, [math.QA/0206223](#).
- [10] J. Hurtubise and E. Markman, Elliptic Sklyanin integrable systems for arbitrary reductive groups, preprint [math.QA/0203031](#).
- [11] M. Kashiwara, in *Algebraic groups and related topics (Kyoto/Nagoya, 1983)*, ed. by R. Hotta, *Adv. Stud. Pure Math.* **6**, (North-Holland, Amsterdam, 1985), pp. 67–81.

- [12] K. Ueno, In *Vector bundles in algebraic geometry (Durham, 1993)*, ed. by N. J. Hitchin, P. E. Newstead and W. M. Oxbury, *London Math. Soc. Lecture Note Ser.* **208**, (Cambridge Univ. Press, Cambridge, 1995) pp. 283–345,
- [13] Y. Shimizu, K. Ueno, *Moduli theory III*, (Iwanami, Tokyo, 1999) Gendai Suugaku no Tenkai series (in Japanese); *Advances in moduli theory*, (English translation by American Mathematical Society, Providence, U.S.A. 2002) Translations of Mathematical Monographs, **206**, Iwanami Series in Modern Mathematics.
- [14] D. Kazhdan, G. Lusztig, *J. Amer. Math. Soc.* **6**, 949–1011 (1993).